

LONG BEACH

3605 Long Beach Blvd #332, Long Beach Ca 90807

FOUNTAIN VALLEY

18122 Brookhurst #B, Fountain Valley Ca 92708

Call or Text (714) 396-0266




WHEN THEIR LIFE MATTERS
EVERY SECOND COUNTS

○ 2020 Advanced Cardiac Life Support




ADVANCED
CARDIOVASCULAR
LIFE SUPPORT

Adult Basic Life Support Algorithm for Healthcare Providers




Adult Cardiac Arrest Algorithm



CPR Quality
<ul style="list-style-type: none"> • Push hard (at least 2 inches [5 cm]) and fast (100-120/min) and allow complete chest recoil. • Minimize interruptions in compressions. • Avoid excessive ventilation. • Change compressor every 2 minutes, or sooner if fatigued. • If no advanced airway, 30:2 compression-ventilation ratio, or 1 breath every 6 seconds. • Quantitative waveform capnography <ul style="list-style-type: none"> - If PETCO₂ is low or decreasing, reassess CPR quality.
Shock Energy for Defibrillation
<ul style="list-style-type: none"> • Biphasic: Manufacturer recommendation (eg, initial dose of 120-200 J); if unknown, use maximum available. Second and subsequent doses should be equivalent, and higher doses may be considered. • Monophasic: 360 J
Drug Therapy
<ul style="list-style-type: none"> • Epinephrine IV/IO dose: 1 mg every 3-5 minutes • Amiodarone IV/IO dose: First dose: 300 mg bolus. Second dose: 150 mg. or • Lidocaine IV/IO dose: First dose: 1-1.5 mg/kg. Second dose: 0.5-0.75 mg/kg.
Advanced Airway
<ul style="list-style-type: none"> • Endotracheal intubation or supraglottic advanced airway • Waveform capnography or capnometry to confirm and monitor ET tube placement • Once advanced airway in place, give 1 breath every 6 seconds (10 breaths/min) with continuous chest compressions
Return of Spontaneous Circulation (ROSC)
<ul style="list-style-type: none"> • Pulse and blood pressure • Abrupt sustained increase in PETCO₂ (typically ≥40 mm Hg) • Spontaneous arterial pressure waves with intra-arterial monitoring
Reversible Causes
<ul style="list-style-type: none"> • Hypovolemia • Hypoxia • Hydrogen ion (acidosis) • Hypo-/hyperkalemia • Hypothermia • Tension pneumothorax • Tamponade, cardiac • Toxins • Thrombosis, pulmonary • Thrombosis, coronary

© 2020 American Heart Association

Adult Cardiac Arrest Circular Algorithm



CPR Quality

- Push hard (at least 2 inches [5 cm]) and fast (100-120/min) and allow complete chest recoil.
- Minimize interruptions in compressions.
- Avoid excessive ventilation.
- Change compressor every 2 minutes, or sooner if fatigued.
- If no advanced airway, 30:2 compression-ventilation ratio.
- Quantitative waveform capnography
 - If PETCO₂ is low or decreasing, reassess CPR quality.

Shock Energy for Defibrillation

- **Biphasic:** Manufacturer recommendation (eg, initial dose of 120-200 J); if unknown, use maximum available. Second and subsequent doses should be equivalent, and higher doses may be considered.
- **Monophasic:** 360 J

Drug Therapy

- **Epinephrine IV/IO dose:** 1 mg every 3-5 minutes
- **Amiodarone IV/IO dose:** First dose: 300 mg bolus. Second dose: 150 mg.
- or
- **Lidocaine IV/IO dose:** First dose: 1-1.5 mg/kg. Second dose: 0.5-0.75 mg/kg.

Advanced Airway

- Endotracheal intubation or supraglottic advanced airway
- Waveform capnography or capnometry to confirm and monitor ET tube placement
- Once advanced airway in place, give 1 breath every 6 seconds (10 breaths/min) with continuous chest compressions


Return of Spontaneous Circulation (ROSC)

- Pulse and blood pressure
- Abrupt sustained increase in PETCO₂ (typically ≥40 mm Hg)
- Spontaneous arterial pressure waves with intra-arterial monitoring


Reversible Causes

- | | |
|---------------------------|-------------------------|
| • Hypovolemia | • Tension pneumothorax |
| • Hypoxia | • Tamponade, cardiac |
| • Hydrogen ion (acidosis) | • Toxins |
| • Hypo-/hyperkalemia | • Thrombosis, pulmonary |
| • Hypothermia | • Thrombosis, coronary |


BLS Termination of Resuscitation




© 2020 American Heart Association



Adult Bradycardia Algorithm




Opioid-Associated Emergency for Lay Responders Algorithm




*For adult and adolescent victims, responders should perform compressions and rescue breaths for opioid-associated emergencies if they are trained and perform Hands-Only CPR if not trained to perform rescue breaths. For infants and children, CPR should include compressions with rescue breaths.

© 2020 American Heart Association

Adult Tachycardia With a Pulse Algorithm



**ACLS Healthcare Provider
Post-Cardiac Arrest Care Algorithm**



Initial Stabilization Phase

Resuscitation is ongoing during the post-ROSC phase, and many of these activities can occur concurrently. However, if prioritization is necessary, follow these steps:

- Airway management: Waveform capnography or capnometry to confirm and monitor endotracheal tube placement
- Manage respiratory parameters: Titrate FiO_2 for SpO_2 92%-98%; start at 10 breaths/min; titrate to $PaCO_2$ of 35-45 mm Hg
- Manage hemodynamic parameters: Administer crystalloid and/or vasopressor or inotrope for goal systolic blood pressure >90 mm Hg or mean arterial pressure >65 mm Hg

Continued Management and Additional Emergent Activities


These evaluations should be done concurrently so that decisions on targeted temperature management (TTM) receive high priority as cardiac interventions.

- Emergent cardiac intervention: Early evaluation of 12-lead electrocardiogram (ECG); consider hemodynamics for decision on cardiac intervention
- TTM: If patient is not following commands, start TTM as soon as possible; begin at 32-36°C for 24 hours by using a cooling device with feedback loop
- Other critical care management
 - Continuously monitor core temperature (esophageal, rectal, bladder)
 - Maintain normoxia, normocapnia, euglycemia
 - Provide continuous or intermittent electroencephalogram (EEG) monitoring
 - Provide lung-protective ventilation


H's and T's

- Hypovolemia**
- Hypoxia**
- Hydrogen ion (acidosis)**
- Hypokalemia/hyperkalemia**
- Hypothermia**
- Tension pneumothorax**
- Tamponade, cardiac**
- Toxins**
- Thrombosis, pulmonary**
- Thrombosis, coronary**

Cardiac Arrest in Pregnancy In-Hospital ACLS Algorithm



Maternal Cardiac Arrest
<ul style="list-style-type: none"> • Team planning should be done in collaboration with the obstetric, neonatal, emergency, anesthesiology, intensive care, and cardiac arrest services. • Priorities for pregnant women in cardiac arrest should include provision of high-quality CPR and relief of aortocaval compression with lateral uterine displacement. • The goal of perimortem cesarean delivery is to improve maternal and fetal outcomes. • Ideally, perform perimortem cesarean delivery in 5 minutes, depending on provider resources and skill sets.
Advanced Airway
<ul style="list-style-type: none"> • In pregnancy, a difficult airway is common. Use the most experienced provider. • Provide endotracheal intubation or supraglottic advanced airway. • Perform waveform capnography or capnometry to confirm and monitor ET tube placement. • Once advanced airway is in place, give 1 breath every 6 seconds (10 breaths/min) with continuous chest compressions.
Potential Etiology of Maternal Cardiac Arrest
<ul style="list-style-type: none"> A Anesthetic complications B Bleeding C Cardiovascular D Drugs E Embolic F Fever G General nonobstetric causes of cardiac arrest (H's and T's) H Hypertension



*Frequent experience or frequent retraining is recommended for providers who perform endotracheal intubation.

Reduce provider exposure

- Don PPE before entering the room/scene
- Limit personnel
- Consider using mechanical CPR devices for adults and adolescents who meet height and weight criteria
- Communicate COVID-19 status to any new providers

Prioritize oxygenation and ventilation strategies with lower aerosolization risk


- Use a HEPA filter, if available, for all ventilation
- Intubate early with a cuffed tube, if possible, and connect to mechanical ventilator, when able
- Engage the intubator with highest chance of first-pass success
- Pause chest compressions to intubate
- Consider use of video laryngoscopy, if available
- Before intubation, use a bag-mask device (or T-piece in neonates) with a HEPA filter and a tight seal
- For adults, consider passive oxygenation with nonrebreathing face mask as alternative to bag-mask device for short duration
- If intubation delayed, consider supraglottic airway
- Minimize closed circuit disconnections

Consider resuscitation appropriateness

- Address goals of care
- Adopt policies to guide determination, taking into account patient risk factors for survival


BLS Healthcare Provider Adult Cardiac Arrest Algorithm for Suspected or Confirmed COVID-19 Patients

Updated April 2020



ACLS Cardiac Arrest Algorithm for Suspected or Confirmed COVID-19 Patients

Updated April 2020



© 2020 American Heart Association

CPR Quality

- Push hard (at least 2 inches [5 cm]) and fast (100-120/min) and allow complete chest recoil.
- Minimize interruptions in compressions.
- Avoid excessive ventilation.
- Change compressor every 2 minutes, or sooner if fatigued.
- If no advanced airway, 30:2 compression-ventilation ratio.
- Quantitative waveform capnography
 - If PETCO₂ <10 mm Hg, attempt to improve CPR quality.
- Intra-arterial pressure
 - If relaxation phase (diastolic) pressure <20 mm Hg, attempt to improve CPR quality.

Shock Energy for Defibrillation

- Biphasic:** Manufacturer recommendation (eg, initial dose of 120-200 J); if unknown, use maximum available. Second and subsequent doses should be equivalent, and higher doses may be considered.
- Monophasic:** 360 J

Advanced Airway

- Minimize closed-circuit disconnection
- Use intubator with highest likelihood of first pass success
- Consider video laryngoscopy
- Endotracheal intubation or supraglottic advanced airway
- Waveform capnography or capnometry to confirm and monitor ET tube placement
- Once advanced airway in place, give 1 breath every 6 seconds (10 breaths/min) with continuous chest compressions

Drug Therapy

- Epinephrine IV/IO dose:** 1 mg every 3-5 minutes
- Amiodarone IV/IO dose:** First dose: 300 mg bolus. Second dose: 150 mg.
or
Lidocaine IV/IO dose: First dose: 1-1.5 mg/kg. Second dose: 0.5-0.75 mg/kg.

Return of Spontaneous Circulation (ROSC)


- Pulse and blood pressure
- Abrupt sustained increase in PETCO₂ (typically ≥40 mm Hg)
- Spontaneous arterial pressure waves with intra-arterial monitoring

Reversible Causes

- Hypovolemia
- Hypoxia
- Hydrogen ion (acidosis)
- Hypo-/hyperkalemia
- Hypothermia
- Tension pneumothorax
- Tamponade, cardiac
- Toxins
- Thrombosis, pulmonary
- Thrombosis, coronary


**BLS Healthcare Provider
Pediatric Cardiac Arrest Algorithm for 2 or More Rescuers
for Suspected or Confirmed COVID-19 Patients**

Updated April 2020




**BLS Healthcare Provider
Pediatric Cardiac Arrest Algorithm for the Single Rescuer
for Suspected or Confirmed COVID-19 Patients**

Updated April 2020



Pediatric Cardiac Arrest Algorithm for Suspected or Confirmed COVID-19 Patients

Updated April 2020



© 2020 American Heart Association

CPR Quality
<ul style="list-style-type: none"> • Push hard ($\geq\frac{1}{3}$ of anteroposterior diameter of chest) and fast (100-120/min) and allow complete chest recoil. • Minimize interruptions in compressions. • Avoid excessive ventilation. • Change compressor every 2 minutes, or sooner if fatigued. • If no advanced airway, 15:2 compression-ventilation ratio.
Shock Energy for Defibrillation
<p>First shock 2 J/kg, second shock 4 J/kg, subsequent shocks ≥ 4 J/kg, maximum 10 J/kg or adult dose</p>
Advanced Airway
<ul style="list-style-type: none"> • Minimize closed-circuit disconnection • Use intubator with highest likelihood of first pass success • Consider video laryngoscopy • Prefer cuffed endotracheal tube if available • Endotracheal intubation or supraglottic advanced airway • Waveform capnography or capnometry to confirm and monitor ET tube placement • Once advanced airway in place, give 1 breath every 6 seconds (10 breaths/min) with continuous chest compressions
Drug Therapy
<ul style="list-style-type: none"> • Epinephrine IO/IV dose: 0.01 mg/kg (0.1 mL/kg of the 0.1 mg/mL concentration). Repeat every 3-5 minutes. • Amiodarone IO/IV dose: 5 mg/kg bolus during cardiac arrest. May repeat up to 2 times for refractory VF/pulseless VT. or • Lidocaine IO/IV dose: Initial: 1 mg/kg loading dose. Maintenance: 20-50 mcg/kg per minute infusion (repeat bolus dose if infusion initiated >15 minutes after initial bolus therapy).
Return of Spontaneous Circulation (ROSC)
<ul style="list-style-type: none"> • Pulse and blood pressure • Spontaneous arterial pressure waves with intra-arterial monitoring
Reversible Causes
<ul style="list-style-type: none"> • Hypovolemia • Hypoxia • Hydrogen ion (acidosis) • Hypoglycemia • Hypo-/hyperkalemia • Hypothermia • Tension pneumothorax • Tamponade, cardiac • Toxins • Thrombosis, pulmonary • Thrombosis, coronary

LONG BEACH

3605 Long Beach Blvd #332, Long Beach Ca 90807

FOUNTAIN VALLEY


18122 Brookhurst #B, Fountain Valley Ca 92708

Call or Text (714) 396-0266



WHEN THEIR LIFE MATTERS
EVERY SECOND COUNTS

○ 2020 Advanced Cardiac Life Support



ADVANCED
CARDIOVASCULAR
LIFE SUPPORT